

Hard m_t Corrections as a Probe of the Symmetry Breaking Sector

J. Bernabéu, D. Comelli, A. Pich and A. Santamaria

Departament de Física Teòrica, IFIC,
CSIC-Universitat de València
46100 Burjassot, València, Spain

Abstract

Non-decoupling effects related to a large m_t affecting non-oblique radiative corrections in vertices ($Z\bar{b}b$) and boxes ($B-\bar{B}$ mixing and ϵ_K) are very sensitive to the particular mechanism of spontaneous symmetry breaking. We analyze these corrections in the framework of a chiral electroweak standard model and find that there is only one operator in the effective lagrangian which modifies the longitudinal part of the W^+ boson without touching the oblique corrections. The inclusion of this operator affects the $Z\bar{b}b$ vertex, the $B-\bar{B}$ mixing and the CP-violating parameter ϵ_K , generating interesting correlations among the hard $m_t^4 \log m_t^2$ corrections to these observables, for example, the maximum vertex $Z\bar{b}b$ correction allowed by low energy physics is about one percent.

One of the basic ingredients of the standard model (SM) is the spontaneous breaking of the electroweak gauge symmetry. In the SM it is implemented through the Higgs mechanism in which the would-be Goldstone excitations are absorbed into the longitudinal degrees of freedom of the gauge bosons. The spontaneous symmetry breaking (SSB) is realized linearly, that means, by the use of a scalar field which acquires a non-zero vacuum expectation value. The spectrum of physical particles contains then not only the massive vector bosons but also a neutral scalar Higgs field which must be relatively light.

In a more general scenario, the SSB can be parametrized in terms of a non-renormalizable lagrangian which contains the SM gauge symmetry realized non-linearly^{1,2}. This non-linearly realized SM is also called the chiral realization of the SM (χ SM) due to of its similarity with low energy QCD chiral lagrangians. It includes, with a particular choice of the parameters of the lagrangian, the SM, as long as the energies involved are small compared with the Higgs mass which is not present in the effective Lagrangian. In addition it can also accommodate any model that reduces to the SM at low energies as happens in many technicolour scenarios. The price to be payed for this general parametrization is the lose of renormalizability and, therefore, the appearance of many couplings which must be determined from experiment or computed in a more fundamental theory.

Since the SSB is related to the bosonic sector, one would expect that any deviation from the SM SSB mechanism would affect especially the gauge boson propagation properties, the so-called oblique corrections, which are parametrized in terms of the S,T,U parameters³ (or the ϵ_1, ϵ_2 and ϵ_3 parameters⁴). In fact these corrections have been extensively studied in the framework of the χ SM⁵. In particular, one would think that one should look into quantities which are M_H -dependent in the SM to test the SSB sector. However, it is interesting to realize that the only M_H -dependent radiative correction, $\Delta\rho$, has an agreement with the SM prediction at the per-mil level. Vertex corrections, whose M_H dependence appears only at the two loop level, are not so well known*.

On the other hand, the would-be Goldstone bosons coming from SSB also couple to fermions. In fact, all non-decoupling effects of the SM related to a large top quark mass, m_t , come from the coupling of the would-be Goldstone bosons to the top quark. Therefore, we can expect any non-decoupling quantity related to a heavy top-quark to be sensitive to the would-be Goldstone boson propagation properties and couplings, that is, to the specific mechanism of SSB.

In the SM, large m_t^2 effects appear, in addition to the oblique corrections, in the vertex $Zb\bar{b}$, that is in $R_b = \Gamma_b/\Gamma_h$, and in $B-\bar{B}$ and $K-\bar{K}$ mixing[†]. Then, we will use these quantities to explore possible deviations of the SM spontaneous symmetry breaking mechanism. To do so we will use a χ SM only for the bosonic sector of the theory and leave fermion couplings as in the linear SM.

It turns out that there is only one operator in the effective lagrangian that affects the $Zb\bar{b}$ vertex without touching the oblique corrections (which as commented before agree with the SM at the per-mil level). This operator modifies the propagation properties of the charged would-be Goldstone bosons, that is, the longitudinal component of the W^+ boson. Therefore, it will also affect any observable in which the non-decoupling effects of a large m_t are important, in particular $B-\bar{B}$

*And, in fact, in the last years there has been a big controversy about the $R_b = \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{hadrons})$ value.

[†]Of course, non-decoupling effects appear in other observables, but only in the quantities we just mentioned present experiments are sensitive enough to see the effects

mixing, and ϵ_K .

In the non-linear realization of the SM the Goldstone bosons π^a associated to the SSB of $SU(2)_L \times SU(2)_R \rightarrow SU(2)_{L+R}$ are collected in a matrix field $U(x) = \text{Exp}\{i\pi^a\tau^a/v\}$. The operators in the effective chiral Lagrangian are classified according to the number of covariant derivatives acting on $U(x)$.

The lowest-order operators just fix the values of the Z and W mass at tree level and do not carry any information on the underlying physics. Therefore, in order to extract some information on new physics we must start studying the effects coming from higher-order operators. Departure of those coefficients from the SM predictions can be a hint for the existence of new physics.

The lowest order effective chiral lagrangian can be written in the following way:

$$L = L_B + L_\psi + L_Y , \quad (1)$$

where

$$L_B = -\frac{1}{2}\text{Tr}\{\hat{W}_{\mu\nu}\hat{W}^{\mu\nu} + \hat{B}_{\mu\nu}\hat{B}^{\mu\nu}\} + \frac{v^2}{4}\text{Tr}\{D_\mu U^\dagger D^\mu U\} , \quad (2)$$

with $\hat{W}_{\mu\nu} = W_{\mu\nu}^a\tau^a/2$, $\hat{B}_{\mu\nu} = B_{\mu\nu}\tau^3/2$, and $D^\mu U = \partial^\mu U + i\frac{g}{2}W_a^\mu\tau^a U - i\frac{g'}{2}B^\mu U\tau^3$. L_ψ is the usual fermionic kinetic lagrangian and

$$L_Y = -\bar{Q}_L U M_q Q_R + h.c , \quad (3)$$

where M_q is a 2×2 block-diagonal matrix containing the 3×3 mass matrices of the up and down quarks and Q_L and Q_R are doublets containing the up and down quarks for the three families in the weak basis.

At the next order, that is containing at most four derivatives, the CP and $SU(2)_L \otimes U(1)_Y$ invariant effective chiral Lagrangian with only gauge bosons and Goldstone fields, is described by the 15 operators reported in ref. 2: $L = \sum_{i=0}^{14} a_i O_i$.

The usual oblique corrections are only sensitive to a_0 , $(a_8 + a_{13})$ and $(a_1 + a_{13})$. On the other hand, the operators proportional to a_2 , a_3 , a_9 and a_{14} parametrize the effective non-abelian gauge couplings that are tested by LEP2. All the other couplings remain not tested because they only contribute to four-point Green functions ($a_4, a_5, a_6, a_7, a_{10}$) or because, although quadratic in the Goldstone fields, they do not contribute to the one-loop oblique corrections (a_{11}, a_{12}). For instance, the operator proportional to a_{11} :

$$O_{11} = \text{Tr}\{(D_\mu V^\mu)^2\} , \quad (4)$$

with $V_\mu = (D_\mu U)U^\dagger$ and $D^\mu V_\mu = \partial^\mu V_\mu + ig[\hat{W}_\mu, V^\mu]$ generates corrections to the two point Green function of the $W^\pm$, Z and would-be Goldstone bosons:

$$\begin{aligned} O_{11} = & g^2 W_\mu^+ \partial^\mu \partial^\nu W_\nu^- + \frac{g_Z^2}{2} Z_\mu \partial^\mu \partial^\nu Z_\nu - 4\pi^+ \frac{\partial^4}{v^2} \pi^- \\ & - 2\pi_3 \frac{\partial^4}{v^2} \pi_3 + \frac{2g}{v} W_\mu^+ \partial^\mu \partial^2 \pi^- + \frac{2g}{v} W_\mu^- \partial^\mu \partial^2 \pi^+ + \frac{2g_Z}{v} Z_\mu^+ \partial^\mu \partial^2 \pi_3 + O(\phi^3) . \end{aligned} \quad (5)$$

However, all these interactions involve always the longitudinal components of the gauge bosons and so do not enter directly into the ϵ_i parameters. The same happens to the operators O_{12} and O_{13} which affect only the longitudinal part of the neutral Z boson.

The effects of the operator O_{11} can be seen more easily once we use the following equation of motion involving the operators of the Lagrangian to lowest order [‡]:

$$D_\mu V^\mu = \frac{i}{v^2} D_\mu \left(\bar{Q}_L \gamma^\mu \tau^a Q_L \tau^a \right) , \quad (6)$$

$$i \not{D} Q_L = U M_q Q_R , \quad i \not{D} Q_R = M_q^+ U^+ Q_L . \quad (7)$$

Then the operator O_{11} can be rewritten as

$$O_{11} = \frac{g^4}{8M_W^4} \left[\bar{Q} (\tau^a U M_q P_R - M_q^+ U^+ \tau^a P_L) Q \right]^2 , \quad (8)$$

where P_L and P_R are the left and right chirality projectors. By writing (8) in terms of the mass eigenstates and keeping only the terms proportional to the top quark mass we obtain

$$O_{11} = \frac{g^4}{8M_W^4} m_t^2 \left((\bar{t} \gamma_5 t)^2 - 4 \sum_{f,f'}^{d,s,b} (\bar{f}'_L t_R) (\bar{t}_R f_L) V_{tf} V_{tf'}^* \right) . \quad (9)$$

Therefore, the effect to lowest order of the modification of the would-be Goldstone propagator can be written as a four-fermion interaction proportional to quark masses. This kind of operators appears also in the analysis of new physics with an effective Lagrangian with SSB realized linearly⁶.

Four fermion interactions are much more convenient for explicit calculations and also to understand the effects of the new operator. For instance, it is clear that the four-fermion interaction can only contribute to the gauge-boson self-energies at two loops and therefore do not contribute to the ϵ_i parameters at one loop.

We discuss now some observables affected by the new interaction.

R_b .—

We start with the evaluation of the corrections to the $Z \bar{b} b$ vertex. We parametrize the effective $Z \bar{b} b$ vertex as:

$$\frac{g}{c_W} Z^\mu \left(g_L^b \bar{b}_L \gamma_\mu b_L + g_R^b \bar{b}_R \gamma_\mu b_R \right) , \quad (10)$$

with the values of the tree level couplings, $g_L^b = -1/2 + s_W^2/3$ and $g_R^b = s_W^2/3$.

At one loop we parametrize the effect of new physics as a shift in the couplings:

$$g_{L,R}^b \rightarrow g_{L,R}^b + \delta g_{L,R}^b . \quad (11)$$

We calculate the one-loop contribution of the operator O_{11} keeping only the divergent logarithmic piece. This means we neglect any possible local contribution from the chiral lagrangian at order

[‡]This is allowed in the effective Lagrangian, even at the one loop level, as long as we keep only the dominant pieces. The use of the equations of motion is equivalent to a redefinition of the fields which affects only higher order operators in the effective Lagrangian

p^6 . The relevant diagram[§] is depicted in fig. (1.a) and the result is:

$$\delta g_L = -\frac{\alpha}{4\pi s_w^2} a_{11} \frac{g^2}{4} \frac{m_t^4}{M_W^4} \log \frac{\Lambda^2}{m_t^2} . \quad (12)$$

A shift in the $Zb\bar{b}$ couplings gives a shift in R_b given by

$$R_b = R_b^{SM} \frac{1 + \delta_{bV}^{NP}}{1 + R_b^{SM} \delta_{bV}^{NP}} , \quad (13)$$

with

$$\delta_{bV}^{NP} = \frac{\delta \Gamma_b}{\Gamma_b^{SM}} \approx 2 \frac{g_L^b}{(g_L^b)^2 + (g_R^b)^2} \delta g_L^b = -4.58 \delta g_L^b . \quad (14)$$

The ALEPH collaboration has presented a new analysis of R_b data which leads to results which are compatible with the standard model predictions at the one sigma level⁷. In fact the new world average is⁸ $R_b = 0.2178 \pm 0.0011$ to be compared with the SM expectation for $m_t = 175$ GeV $R_b^{SM} = 0.2157 \pm 0.0002$. Clearly the new value of R_b is within two standard deviations of the standard model predictions⁹.

Using these data on R_b we get

$$\delta_{bV}^{NP} = 0.012 \pm 0.007 . \quad (15)$$

$K-\bar{K}$ and $B-\bar{B}$ Mixing .–

In the SM, the mixing between the B^0 meson and its antiparticle is completely dominated by the top contribution. The explicit m_t dependence of the corresponding box diagram is given by the loop function¹⁰

$$S(x_t)_{SM} = \frac{x_t}{4} \left[1 + \frac{9}{1-x_t} - \frac{6}{(1-x_t)^2} - \frac{6x_t^2 \ln x_t}{(1-x_t)^3} \right] , \quad x_t \equiv \frac{\overline{m}_t^2}{M_W^2} , \quad (16)$$

which contains the hard m_t^2 term, $S(x_t) \sim x_t/4$, induced by the longitudinal W exchanges. The same function regulates the top-quark contribution to the $K-\bar{K}$ mixing parameter ε_K . The measured top-mass, $m_t = 175 \pm 6$ GeV [$\overline{m}_t \equiv \overline{m}_t(m_t) = 167 \pm 6$ GeV], implies $S(x_t)_{SM} = 2.40 \pm 0.13$.

The correction induced by the new operator, O_{11} , can be parametrized as a shift on the function $S(x_t)$. The calculation of the diagrams[§] in fig. (1.b) leads to the result:

$$S(x_t) = S(x_t)_{SM} + \delta S(x_t) , \quad \delta S(x_t) = -a_{11} \frac{g^2 m_t^4}{2M_W^4} \ln \frac{\Lambda^2}{m_t^2} . \quad (17)$$

[§]The same result is of course obtained using the original form (4) for O_{11} , where the effect of this operator appears as a modification of the longitudinal W propagator. However, one needs to consider a larger number of Feynman diagrams in this case.

Thus, the hard $m_t^4 \ln m_t^2$ contributions to δ_{bV}^{NP} and $\delta S(x_t)$ are correlated:

$$\delta S(x_t) = \frac{32\pi^2}{|V_{tb}|^2 g^2} \delta g_L^b = -163 \delta_{bV}^{NP}. \quad (18)$$

We can use the measured $B_d^0-\bar{B}_d^0$ mixing¹¹, $\Delta M_{B_d^0} = (0.464 \pm 0.018) \times 10^{12} \text{ s}^{-1}$, to infer the experimental value of $S(x_t)$ and, therefore, to set a limit on the δg_L^b contribution. The explicit dependence on the quark-mixing parameters can be resolved putting together the constraints from $\Delta M_{B_d^0}$, ε_K and $\Gamma(b \rightarrow u)/\Gamma(b \rightarrow c)$. Using the Wolfenstein parametrization¹² of the quark-mixing matrix, one has:

$$\left| \frac{V_{td}}{\lambda V_{cb}} \right| = \sqrt{(1-\rho)^2 + \eta^2} = \frac{(1.21 \pm 0.09)}{\sqrt{S(x_t)}} \frac{185 \text{ MeV}}{\sqrt{\eta_B} (\sqrt{2} f_B \sqrt{B_B})} = \frac{(1.21_{-0.30}^{+0.50})}{\sqrt{S(x_t)}}, \quad (19)$$

$$\eta \left[(1-\rho) A^2 \eta_2 S(x_t) + P_0 \right] A^2 B_K = 0.226 \quad (20)$$

$$\left| \frac{V_{ub}}{\lambda V_{cb}} \right| = \sqrt{\rho^2 + \eta^2} = 0.36 \pm 0.09. \quad (21)$$

We have taken $\lambda \equiv |V_{us}| = 0.2205 \pm 0.0018$, $|V_{cb}| \equiv A\lambda^2 = 0.040 \pm 0.003$ and $|V_{ub}|/|V_{cb}| = 0.08 \pm 0.02$. The numerical factor on the rhs of eq. (19) should be understood as an allowed range, because the error is dominated by the large theoretical uncertainties in the hadronic matrix element of the $\Delta B = 2$ operator; it corresponds to^{13,14} $\sqrt{\eta_B} (\sqrt{2} f_B \sqrt{B_B}) = (185 \pm 45) \text{ MeV}$. In eq. (20), $\eta_2 = 0.57 \pm 0.01$ is the short-distance QCD correction¹⁵, while $P_0 = 0.31 \pm 0.02$ takes into account the charm contributions¹³. For the $\Delta S = 2$ hadronic matrix element we have chosen the range¹⁴ $B_K = 0.6 \pm 0.2$.

Both the circle (19) and the hyperbola (20) depend on the value of $S(x_t)$. The intersection of the two circles (19) and (21) restricts $S(x_t)$ to be in the range $0.39 < |S(x_t)| < 9.7$. The request of simultaneous intersection with the hyperbola ϵ_K imposes a further constraint. Since a positive value of B_K is obtained by all present calculations and $S(x_t)_{\text{SM}} > 0$, the SM implies a positive value for η . In our case, the constraint that the total $S(x_t) = S(x_t)_{\text{SM}} + \delta S(x_t)$ is positive does not exist and this opens the possibility of solutions also with $\eta < 0$; however, this would imply a huge correction $\delta S(x_t)$. Taking $\eta > 0$, the three curves (bands) intersect if $S(x_t) > S(x_t)_{\text{min}} = 1.0$.

The minimum value of $S(x_t)$ is reached for V_{cd}^{max} , B_K^{max} and $|V_{ub}/V_{cb}|^{\text{max}}$. Taking a more conservative ± 0.14 error in eq. (21) (corresponding to $|V_{ub}/V_{cb}| = 0.08 \pm 0.03$) would result in $S(x_t)_{\text{min}} = 0.8$.

The shift in g_L^b required by R_b [eq. (15)] and the relation (18) imply

$$\delta S = -2.0 \pm 1.1, \quad (22)$$

i.e., $-0.7 < S(x_t) < 1.5$. Thus, the present experimental measurements of R_b and the low-energy constraints from the usual *unitarity triangle fits* are compatible with the introduction of the operator O_{11} . From eq. (18) and eq. (15) and the constraint $S \geq S_{\text{min}} = 1$ we can see that the maximum (positive) value of δ_{bV}^{NP} allowed by low-energy physics is

$$\delta_{bV}^{NP} < 0.01,$$

which is even stronger than the values obtained by present direct measurements of R_b (eq. (15)).

We acknowledge interesting discussions with Misha Bilenky, Domenec Espriu and Joaquim Matias. One of us (D.C.) is indebted to the Spanish Ministry of Education and Science for a postdoctoral fellowship. This work has been supported by the Grant AEN-96/1718 of CICYT, Spain.

References

- [1] T. Appelquist, “Gauge Theories and Experiments at High Energies”, ed. K.C. Brower and D.G. Sutherland (Scottish University Summer School in Physics, St. Andrews); T. Appelquist and C. Bernard, Phys. Rev. **D22**, 200 (1980)
- [2] A.C. Longhitano, Phys. Rev. **D22**, 1166 (1980); Nucl. Phys. **B188**, 188 (1981)
- [3] M.E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990); Phys. Rev. **D46**, 381 (1992); W. Marciano and J. Rosner, Phys. Rev. Lett **65**, 2963 (1990)
- [4] G. Altarelli and R. Barbieri, Phys. Lett. **B253**, 161 (1990); G. Altarelli, R. Barbieri and J. Jadach, Nucl. Phys. **B369**, 3 (1992)
- [5] See for instance F. Feruglio, Int. J. Mod. Phys. **A8**, 4937 (1993)
- [6] J. Bernabeu, D. Comelli, A. Pich and A. Santamaria in preparation.
- [7] I. Tomalin, “ R_b with Multivariate Analysis”, talk at the ICHEP 96 Conference, Warsaw, July 1996.
- [8] The LEP Electroweak Working Group and the SLD Heavy Flavor Group, “A Combination of Preliminary LEP and SLD Electroweak Measurements and Constraints on the Standard Model”. Report LEPEWWG/96-02, 1996.
- [9] A. Akundov, D. Bardin and T. Riemann, Nucl. Phys. **B276**, 1 (1986); J. Bernabéu, A. Pich, and A. Santamaria, Phys. Lett. **B200**, 569 (1988); Nucl. Phys. **B363**, 326 (1991); W. Beenakker and W. Hollik, Z. Phys. **C40**, 141 (1988)
- [10] G. Buchalla, A.J. Buras and M.E. Lautenbacher, “Weak Decays Beyond Leading Logarithms” Submitted to Rev. Mod. Phys. [hep-ph/9512380].
- [11] L. Gibbons, *Quark Masses and Mixing from Weak Processes*, Plenary talk at the 28th International Conference on High Energy Physics (Warsaw, 1996)
- [12] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983)

- [13] A.J. Buras, *Flavour Changing Neutral Current Processes*, Plenary talk at the 28th International Conference on High Energy Physics (Warsaw, 1996), [hep-ph/9610461].
- [14] A. Pich and J. Prades, Phys. Lett. **B346**, 342 (1995);
- [15] A.J. Buras, M. Jamin and P.H. Weisz, Nucl. Phys. **B347**, 491 (1990)





Figure 1: a) Contribution of the effective operator O_{11} to $Z \rightarrow b\bar{b}$. b) Contribution of the effective operator O_{11} to B - $\bar{B}$ mixing